

## เอกสารอ้างอิง (References)

- American Diabetes Association. Standards of medical care in diabetes–2007. *Diabetes Care* 2007; 30(Suppl 1): S4-S41.
- Chan, A. Y. W., Swaminathan, R., and Cockram, C. S. (1989). Effective of sodium fluoride as a preservative of glucose in blood. *Clinical Chemistry*, 35(2), 315-317.
- Horowitz L, G. (2008). Reference intervals: practical aspects. *Electronic Journal International Federation of Clinical Chemistry and Laboratory Medicine*, 19(2), 1-11.
- Landt, M. (2000). Glyceraldehyde preserves glucose concentration in whole blood specimens. *Clinical Chemistry*, 46(8), 1144-1149.
- Lolekha, P.H., Vanavanan, S., Lolekha, S. (2001) Update on value of the anion gap in clinical diagnosis and laboratory evaluation. *Clin Chim Acta*, 2001; 307(1-2):33-6.
- O’Keane, M.P. and Sean K. Cunningham, K.S. (2006). Evaluation of three different specimen types (serum, plasma lithium heparin and serum gel separator) for analysis of certain analytes: clinical significance of differences in results and efficiency in use. *Clinical Chemistry Laboratory Medicine*, 44(5):662–668.
- Shi, R.Z., Seeley, E.S., Bowen, R., and Faix, J. (2009). Rapid blood separation is superior to fluoride for preventing in vitro reductions in measured blood glucose concentration. *Journal of Clinical Pathology*, 62(8), 752 - 753.
- Young, D.S. & Bermes E.W. Jr. (1994). Specimen collection and processing; sources of biological variation (2<sup>nd</sup> ed.). In: Burtis C.A., Ashwood E.R., eds. *Tietz textbook of clinical chemistry*. Philadelphia: WB Saunders, 60 - 61.