

Bibliography

- Cai, Y., & Corke, H. (2000). Production and Properties of Spray-dried Amaranthus Betacyanin Pigments. *Journal of food science*, 65(7), 1248-1252.
- Chang, C.-C., Yang, M.-H., Wen, H.-M., & Chern, J.-C. (2002). Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of food and drug analysis*, 10(3).
- Chau, C.-F., Wang, Y.-T., & Wen, Y.-L. (2007). Different micronization methods significantly improve the functionality of carrot insoluble fibre. *Food Chemistry*, 100(4), 1402-1408.
doi:<http://dx.doi.org/10.1016/j.foodchem.2005.11.034>
- Chuacharoen, T., & Sabliov, C. M. (2016). The potential of zein nanoparticles to protect entrapped β -carotene in the presence of milk under simulated gastrointestinal (GI) conditions. *LWT - Food Science and Technology*, 72, 302-309. doi:<http://dx.doi.org/10.1016/j.lwt.2016.05.006>
- Dib Taxi, C., De Menezes, H., Santos, A., & Grosso, C. (2003). Study of the microencapsulation of camu-camu (*Myrciaria dubia*) juice. *Journal of microencapsulation*, 20(4), 443-448.
- Fazaeli, M., Emam-Djomeh, Z., Kalbasi Ashtari, A., & Omid, M. (2012). Effect of spray drying conditions and feed composition on the physical properties of black mulberry juice powder. *Food and Bioproducts Processing*, 90(4), 667-675.
doi:<http://dx.doi.org/10.1016/j.fbp.2012.04.006>
- Footrakul, P., Mawimol, W., & Boonyasupa, W. (2003). *Development of lime powder processing and quality evaluations during storage*. Paper presented at the Proceedings of 41st Kasetsart University Annual Conference, 3-7 February, 2003. Subject: Agro-Industry.
- Ghafar, M., Prasad, K. N., Weng, K. K., & Ismail, A. (2010). Flavonoid, hesperidine, total phenolic contents and antioxidant activities from Citrus species. *African Journal of Biotechnology*, 9(3).
- Gil-Izquierdo, A., Gil, M. I., & Ferreres, F. (2002). Effect of Processing Techniques at Industrial Scale on Orange Juice Antioxidant and Beneficial Health Compounds. *Journal of Agricultural and Food Chemistry*, 50(18), 5107-5114.
doi:10.1021/jf020162+
- Kuljarachanan, T., Devahastin, S., & Chiewchan, N. (2009). Evolution of antioxidant compounds in lime residues during drying. *Food Chemistry*, 113(4), 944-949.

- Largo Avila, E., Cortes Rodríguez, M., & Velásquez, C. (2015). Influence of Maltodextrin and Spray Drying Process Conditions on Sugarcane Juice Powder Quality. *Revista Facultad Nacional de Agronomía, Medellín*, 68(1), 7509-7520.
- Liu, X., Cui, C., Zhao, M., Wang, J., Luo, W., Yang, B., & Jiang, Y. (2008). Identification of phenolics in the fruit of emblica (*Phyllanthus emblica* L.) and their antioxidant activities. *Food Chemistry*, 109(4), 909-915.
- Mara Righetto, A., & Maria Netto, F. (2005). Effect of encapsulating materials on water sorption, glass transition and stability of juice from immature acerola. *International Journal of Food Properties*, 8(2), 337-346.
- Padayatty, S. J., Katz, A., Wang, Y., Eck, P., Kwon, O., Lee, J. H., . . . Levine, M. (2003). Vitamin C as an antioxidant: evaluation of its role in disease prevention. *J Am Coll Nutr*, 22(1), 18-35.
- Quek, S. Y., Chok, N. K., & Swedlund, P. (2007). The physicochemical properties of spray-dried watermelon powders. *Chemical Engineering and Processing: Process Intensification*, 46(5), 386-392.
doi:<http://dx.doi.org/10.1016/j.cep.2006.06.020>
- Singleton, V. L., Orthofer, R., & Lamuela-Raventós, R. M. (1999). [14] Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods in enzymology*, 299, 152-178.
- Shihawong, W. (2003). *Effects of maltodextrin on qualities of lime powder produced by freeze drying*. Paper presented at the Proceedings of 41st Kasetsart University Annual Conference, 3-7 February, 2003. Subject: Agro-Industry.
- Theansuwan, W., Triratanasirichai, K., & Tangchaichit, K. (2008). Continuous production of lime juice by vacuum freeze drying. *American Journal of Applied Sciences*, 5(8), 959-962.
- Tonon, R. V., Brabet, C., & Hubinger, M. D. (2010). Anthocyanin stability and antioxidant activity of spray-dried açai (*Euterpe oleracea* Mart.) juice produced with different carrier agents. *Food Research International*, 43(3), 907-914.
doi:<https://doi.org/10.1016/j.foodres.2009.12.013>